

Secretary Of The Navy Professor

James J. O'Brien
Center For Ocean-Atmospheric Prediction Studies
The Florida State University
Tallahassee, FL 32306-2840
Phone: (850) 644-4581 Fax: (850) 644-4841 Email: obrien@coaps.fsu.edu
Award #: N00014940369
<http://www.coaps.fsu.edu>

LONG-TERM GOAL

The goal of this research is to develop a predictive capability for the upper ocean circulation and atmospheric interactions using numerical models.

OBJECTIVE

The immediate objectives are to develop improved models of the oceans, to evaluate these models against available observations, and to develop assimilation techniques to be used in these models. In addition, we are continuing the task of preparing long-term global surface fluxes for use in ocean modeling.

This special ONR Grant is the base support for the FSU Center for Ocean-Atmospheric Prediction Studies. Other agencies who contribute are NASA, NSF, and NOAA.

APPROACH

We are using a suite of models forced with estimates of real winds, with very fine horizontal resolution and realistic basin geometries. The vertical structure ranges from limited resolution, as in the reduced gravity model, to those with very high vertical resolution, as in the multi-layered models and the Princeton model. Much of the modeling work is done in cooperation with the Naval Research Laboratories in Stennis, MS. We have also been examining surface fluxes for use in large-scale numerical ocean models.

WORK COMPLETED

The circulation of the Yellow, East China and Bohai Seas were modeled using the Princeton Ocean Model to better understand the dynamics of the Yellow Sea Warm Current.

Results from multi-layered numerical models were downloaded and analyzed in the region of Hawaii and the Gulf of Alaska. Model transport, velocity, layer thickness and pressure fields were examined. Calculation of water mass transport were done and model validation was carried out by comparison to observations.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 1999		2. REPORT TYPE		3. DATES COVERED 00-00-1999 to 00-00-1999	
4. TITLE AND SUBTITLE Secretary Of The Navy Professor				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Florida State University,Center For Ocean-Atmospheric Prediction Studies,Tallahassee,FL,32306				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 7	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

We have developed a coupled flux and sea state model that can utilize any one of several wave characteristics to specify the sea state. Variations of the surface temperature in the North Atlantic were analyzed over multi-decadal timescales. Two primary patterns of variation were found.

RESULTS

The Yellow Sea Warm Current was shown to be present throughout the year, reaching its furthest northward extension in winter (c.f. Fig. 1). Sensitivity tests and spectrum analyses show that winds modify the pathway and extent of the Yellow Sea Warm Current. The current's origin, however, appears to be due to external forcing from the current systems developed in the East China Sea.

The North Hawaiian Ridge Current (NHRC, see Fig. 2) was shown to be a robust surface trapped feature of the mean flow in the central Pacific Ocean. The NHRC manifested from a northern branch of the North Equatorial Current (NEC). The NHRC was found to begin at the bifurcation point of the northern branch of the NEC, leaking 2.0 Sv through the Alenuihaha channel to the lee of the islands, and continuing along the Hawaiian Island chain until it reaches Kauai. The NHRC width scale was consistent with that of an inertial boundary layer. The effects of Ekman pumping/suction, vertical mixing, and the meridional overturning cell on the mean flow were negligible. We found abyssal currents steer upper ocean currents resulting in a northward migration of the bifurcation and enhanced transport north of Hawaii. However, the upper ocean-topographic coupling is not evident in the upper layer currents.

We have developed a coupled flux and sea state model that can utilize any one of several wave characteristics to specify the sea state. The input sea state can be specified through wave age, phase speed of the dominant waves, period of the dominant waves, significant wave height, and significant slope (ratio of significant wave height to the wavelength of the dominant waves). Period and significant wave heights are recorded on an increasing number of National Buoy Data Center (NBDC) buoys, and significant wave height is routinely observed with altimeters. These observations, coupled with wind speed, allow us to determine fluxes without arbitrarily specified sea states. The neutral drag coefficient CD_{10N} (Fig. 3) is shown as a function of wind speed and significant wave height. Note the consistency with the observations that CD decreases as wave height increases (for a fixed wind speed) [Willard Pierson, personal communication, 1998].

A 50-year propagating oscillation and 15-year rotating pattern are found in the North Atlantic sea surface temperature anomaly (SST). COADS SST anomaly fields over much of the North Atlantic are investigated using propagating Complex Empirical Orthogonal Functions (CEOFs) during the period of 1947 to 1992. The first mode (58% of the variance) is a northward propagating feature and the second and third modes combined (about 30% of the variance) is a cyclonically rotating feature within the sub-polar gyre.

IMPACTS

Much of this successful project was conducted as a cooperative effort between Naval and academic personnel. Naval operations can use this information to better understand the local environment, enhancing countermeasure measures and deployment.

RELATIONSHIP TO OTHER PROJECTS

COAPS manages an ONR Minority Program that currently places 10-15 minority undergraduate science students in a laboratory-type work position with a mentoring professor. The goal is to convince these students to do graduate studies. Without the basic Secretary of Navy Grant, the ONR Minority Program would not have the infrastructure for the administration.

NSF supports the World Ocean Circulation Experiment Data Assembly Center and Special Analysis Center (WOCE DAC/SAC) for surface meteorology and surface fluxes. This center archives surface meteorological observations from international research vessels as well as develops and validates high-quality surface flux fields, e.g. our tropical Pacific Ocean wind products for El Nino forecasts.

NASA, through a science grant from NASA Headquarters and the NASA Scatterometer Project at JPL, provides support for basic science in utilizing scatterometer data to drive ocean models.




Fig 1. Temperature and currents at 20 m depth from the numerical model of the Yellow Sea. The arrows have been truncated for currents greater than 0.1 m/s.




Fig 2. Model results from the region of Hawaii. The surface currents are represented by the arrows. The layer thickness is represented by color.




Fig 3. Modeled drag coefficient (C_D) as a function of wind speed and a significant wave height. Local wind-wave equilibrium is shown by the green dashed line, rising seas are in red, and falling seas in blue. Conditions that require further validation are shown by the gray background.

PUBLICATIONS

Published

Bourassa, M. A., D. G. Vincent, and W. L. Wood, 1999: A flux parameterization including the effects of capillary waves and sea state, *J. Atmos. Sci.*, Vol. 56, pp. 1123 - 1139.

- Bourassa, M. A., L. Zamudio, and J. J. O'Brien, 1999: Non-inertial flow in NSCAT observations of Tehuantepec winds. *J. Geophys. Res.*, 104 (C5), pp. 11,311 – 11,319.
- Bove, M.C., D.F. Zierden, and J.J. O'Brien, 1998: Are Gulf Hurricanes Getting Stronger? *Bull. Amer. Meteorol. Soc.*, 79, pp. 1327 - 1328.
- Bove, M.C., J.B. Elsner, C.W. Landsea, X. Niu, and J.J. O'Brien, 1998: Effects of El Niño on U.S. Landfalling Hurricanes, Revisited, *Bull. Amer. Meteorol. Soc.*, 79, pp. 2477 - 2482.
- Bulusu, S., I. S. Robinson, J. R. Blundell, and P. G. Challenor, 1999: Indian Ocean Rossby waves observed in TOPEX/POSEIDON altimeter data and in model simulations, *Int. J. of Remote Sensing*, (accepted).
- Caron, J. M., and J. J. O'Brien, 1998: The Generation of Synthetic Sea Surface Temperature Data for the Equatorial Pacific Ocean. *Mon. Wea. Rev.*, 126, No. 11, pp. 2809 - 2821. [Corrigendum: *Mon. Wea. Rev.*, 127, No. 4, p. 575].
- Izaurrealde, R. C., N. J. Rosenberg, R. A. Brown, D. M. Legler, M. Tiscareno-Lopez, R. Srinivasan, and R. D. Sands, 1999: Modeled effects of strong Los Ninos on crop productivity in North America. *Ag. Forest Meteor.*, 94, pp. 259-268.
- Larow, T. E., and T. N. Krishnamurti, 1998: Initial conditions and ENSO prediction using a coupled ocean - atmosphere model. *Tellus*, 50A, pp. 76 - 90.
- Latif, M., D. Anderson, T. Barnett, M. Cane, R. Kleeman, A. Leetmaa, J. O'Brien, A. Rosati, and E. Schneider, 1998: A Review of the Predictability and Prediction of ENSO, *J. Geophys. Res.*, 103 (C7), pp. 14,375 - 14,393.
- Legler, D. M., K. J. Bryant, and J. J. O'Brien, 1999: Impact of ENSO-related climate anomalies on crop yields in the US, *Climatic Change*, 42 (2), pp. 351-375.
- Mask, A. C., J. J. O'Brien, and R. Preller, 1999: Wind driven effects on the Yellow Sea Warm Current, *J. Geophys. Res.*, 103 (C13), pp. 30,713 - 30,729.
- Melsom, A., S. D. Meyers, J. J. O'Brien, H. E. Hurlburt and J. Metzger, 1999: ENSO effects on Gulf of Alaska eddies, *Earth Interac.*, 3.
- Meyers, S. D., and S. K. Basu, 1999: Eddies in the eastern Gulf of Alaska from Topex/Poseidon altimetry, *J. Geophys. Res.*, 104 (C6), pp. 13,333-13,343.
- Meyers, S.D., A. Melsom, G. T. Mitchum, and J.J. O'Brien, 1999: Detection of the fast Kelvin wave teleconnection during ENSO, *J. Geophys. Res.*, 103 (C12), pp. 27,655 - 27,663.
- Meyers, S.D., J.J. O'Brien and E. Thelin, 1999: Reconstruction of monthly SST in the Tropical Pacific Ocean during 1868-1993 using adaptive climate basis functions, *Mon. Wea. Rev.*, 127, pp. 1599-1612.
- Murphy, S.J., H.E. Hurlburt, and J.J. O'Brien, 1999: The connectivity of eddy variability in the Caribbean Sea, the Gulf of Mexico, and the Atlantic Ocean, *J. Geophys. Res. - Oceans*, 104 (C1), pp. 1431 - 1453.
- Rosenberg, N. J., R. C. Izaurrealde, M. Tiscareño - López, D. Legler, R. Srinivasan, R. A. Brown, and R. D. Sands, 1998: Sensitivity of North American Agriculture to ENSO - based Climate Scenarios and their Socio - Economic Consequences: Modeling in an Integrated Assessment Framework. PNL - 11699/UC - 0000, Pacific Northwest National Laboratory, Washington, DC, 146 pp.
- Siefridt, L., B. Barnier, D. M. Legler, and J. J. O'Brien, 1998: 5 - day average wind over North - West Atlantic from ERS1 using a variational analysis, *The Global Atmos. and Ocean Sys.*, 5, pp. 317 - 344.
- Smith, S. R., P. M. Green, A. P. Leonardi, and J. J. O'Brien, 1998: Role of multiple - level tropospheric circulations in forcing ENSO winter precipitation anomalies. *Mon. Wea. Rev.*, *Amer. Meteorol. Soc.*, 126, pp. 3102 - 3116.
- Smith, S. R., M. A. Bourassa, and R. J. Sharp, 1999: Establishing more truth in true winds, *J. Atmos. Oceanic Technol.*, 16, pp. 939-952.

- Solow, A., R. M. Adams, K. J. Bryant, D. M. Legler, J. J. O'Brien, B. A. McCarl, W. Nayda, and R. Weiher, 1998: The value of improved ENSO prediction to US agriculture, *Climatic Change*, 39 (1), pp. 47 - 60.
- Spiesberger, J. L., H. E. Hurlburt, M. A. Johnson, M. Keller, S. D. Meyers and J. J. O'Brien, 1998: Acoustic Thermometry Data Compared With Two Ocean Models: The importance of Rossby Waves and ENSO in Modifying the Ocean Interior, *Dynamics of Atmos. and Oceans.*, 26, 4, pp. 209 - 240.
- Tiscareno Lopez, M., N. J. Rosenberg, D. M. Legler, A. R. Corral, R. Shrinivasan, R. A. Brown, G. G. Medina, M. A. V. Valle, and R. C. Isaurralde, 1998: Algunos efectos del fenomeno climatico El Nino en la agricultura Mexicana. *Ciencia y Desarrollo*, 24, pp. 4 - 14.
- Verschell, M.A., M. A. Bourassa, D. E. Weissman, and J. J. O'Brien, 1999: Ocean Model Validation of the NASA Scatterometer Winds, *J. Geophys. Res.*, 104 (C5), pp. 11,359 – 11,373.
- White, L.D., M. Tewari and T. N. Krishnamurti, 1998: Application of a GCM to study the surface hydrological budget of Amazonia. *J. Appl. Meteor.*, 37, pp. 1321 - 1331.

In Press

- Basu, S. K., S. D. Meyers, and J. J. O'Brien, 1999: Annual and interannual sea level variations in the Indian Ocean from Topex/Poseidon observations and ocean model simulations, *J. Geophys. Res.*, (accepted).
- Bourassa, M. A., 1999: Shear-Stress Model for the Aqueous Boundary-Layer Near the Air-Sea Interface, *J. Geophys. Res. - Oceans*, (in press).
- Cocke, S., and T.E. Larow, 1999: Seasonal Predictions using a Coupled Ocean-Atmosphere Regional Spectral Model. *Mon. Wea. Rev.*, (accepted).
- Cocke, S., and T.E. Larow, 1999: Seasonal Predictions using a Coupled Ocean-Atmosphere Regional Spectral Model. *Mon. Wea. Rev.*, (in press).
- Latif, M., D. Anderson, T. Barnett, M. Cane, R. Kleeman, A. Leetmaa, J. O'Brien, A. Rosati, and E. Schneider, 1999: TOGA Review Paper "Predictability and Prediction", *Mon. Wea. Rev.*, (to appear).
- Meyers, S. D., S. K. Basu, and J. J. O'Brien, 1999: TOPEX/Poseidon altimetry captures cycles of the Indian Ocean, *International WOCE Newsletter*, (to appear).
- Mizoguchi, K., S. D. Meyers, S. K. Basu, and J. J. O'Brien, 1999: Multi- and quasi-decadal variations of sea surface temperature anomaly in the North Atlantic, *J. Phys. Oceanogr.*, (in press).

Submitted

- Bachiochi, D., and T. N. Krishnamurti, 1999: Enhanced Low-Level Stratus in the FSU Coupled Ocean-Atmosphere Model. *J. Clim.*, (submitted).
- Bourassa, M. A., 1999: A Sea State Parameterization with Non-Arbitrary Wave Age Applicable to Low and Moderate Wind Speeds, *J. Phys. Oceanogr.*, (submitted).
- Cocke, S. and T.E. LaRow, 1999: A Nested Coupled Ocean-Atmosphere Model for Climate Studies. *Mon. Wea. Rev.*, (submitted).
- Kreitner, R.E., S.R. Smith, and J.J. O'Brien, 1999: Front range severe flash flooding and El Nino. *Weather and Forecasting*, (submitted).
- Murray, C. P., S. L. Morey, and J. J. O'Brien, 1999: Interannual Variability of Upper Ocean Vorticity Balances in the Gulf of Alaska, *J. Geophys. Res. - Oceans*, (submitted).

- Pegion, P. J., M. A. Bourassa, D. M. Legler, and J. J. O'Brien, 1999: Objectively derived daily pseudostress fields from NSCAT data created through direct minimization, cross validation, and multigridding. *Mon. Wea. Rev.*, (submitted).
- Putman, W. M., D. M. Legler, and J. J. O'Brien, 1999: Synthesis of FSU and NCEP reanalysis pseudostress products over the Pacific ocean. *J. Climate*, (submitted).
- Rao, A. D., D. M. Legler and J. J. O'Brien, 1999: Variability of Surface Curl over the Indian Ocean: 1970-95, *J. Phys. Ocean.*, (submitted).
- Smith, S. R., D. M. Legler, J. J. O'Brien, and M. R. Jarvis, 1999: Comparison of 1997-1998 US Temperature and precipitation to historical ENSO warm phase conditions. *Mon. Wea. Rev.*, (submitted).
- Smith, S.R. and J.J. O'Brien, 1999: Regional snowfall distributions associated with ENSO: Implications for seasonal forecasting. *Weather and Forecasting*, (submitted).
- Verschell, M.A., S.D. Meyers, and J.J. O'Brien, 1999: Interannual variability of atmospheric carbon dioxide flux in the Equatorial Pacific Ocean, *Prog. Ocean.*, (submitted).
- Verzone, K., M. Bourassa, T. E. LaRow, S. D. Cocke, and J. J. O'Brien, 1999: Double Ensemble Estimates of Precipitation in the Southeastern United States for Extreme ENSO Events. *J. Clim.*, (submitted).

Presentations

San Francisco, CA	12-10-97	AGU Meeting
New Orleans, LA	01-09-98	Visit NRL-Stennis
San Diego, CA	02-08-98	AGU Ocean Science Meeting
Philadelphia, PA	02-12-98	AAAS Annual Meeting
Biloxi, MS	04-03-98	ONR Meeting on Univ. and Bus. Technology
Iquique, Chile	05-02-98	Inst. of Oceanography Meeting
St. Louis, MO	06-08-98	NOAA/OGP GCIP Climate Conference
Miami, FL	09-08-98	CIMAS Visiting Scientist Program & RSMAS